



**HELP SPEND YOUR VACATION
WHERE YOU WANT**



AND NOT IN THE BATHROOM.

DUKORAL[®] is a drinkable vaccine that helps prevent diarrhea caused by enterotoxigenic *E. coli* producing heat-labile enterotoxin and cholera.

**ASK YOUR HEALTHCARE PROFESSIONAL ABOUT TAKING
DUKORAL[®] AT LEAST 2 WEEKS BEFORE TRAVELLING.**



*Except in Quebec

Help prevent ETEC diarrhea with
DUKORAL[®]
2 weeks before you leave for 3 months protection

Enterotoxigenic *E. coli* (ETEC) is the most common cause of diarrhea for travellers to many parts of the world.



Enterotoxigenic *E. coli* (ETEC) diarrhea can spread very easily from person to person.



Annually, ETEC is estimated to cause 200 million diarrheal episodes and approximately 380,000 deaths worldwide.



If you eat food or drink water or ice that is contaminated with ETEC, you are at risk of getting ETEC diarrhea.

98% of travellers have been shown to make a food or water mistake within the first 3 days of their vacation:

- Drinking beverages with ice cubes or crushed ice
- Brushing teeth with tap water
- Consuming salads and uncooked vegetables
- Eating fruits which cannot be peeled
- Sampling local cuisine from street vendors

A STUDY SHOWED THAT TRAVELLERS AFFECTED BY ETEC DIARRHEA LOSE AN AVERAGE OF 3.5 DAYS OF VACATION TIME.



ETEC diarrhea can seriously interfere with your travel plans

ETEC diarrhea can last for a few days, keep you in bed, require medical treatment, and result in:

- Nausea
- Vomiting
- Abdominal cramps
- Fever
- Chills
- Loss of appetite
- Headache
- Muscle aches
- Bloating

A study assessed the cost of travellers' diarrhea caused by ETEC

The average cost of an episode of travellers' diarrhea was estimated to be **US \$1,460**:

an average of **3.5 days lost**

X

more than **\$400 per day** in travel, time and medical costs

=

US \$1,460

IT'S YOUR TIME AND MONEY.

Am I at risk?

If you're travelling to a high-risk area, you should consider the threat of ETEC diarrhea.

Incidence of Enterotoxigenic *E. coli* (ETEC)



IN 2011, THE RATES WERE HIGHEST IN LATIN AMERICA/CARIBBEAN, AFRICA AND SOUTH ASIA.

DUKORAL[®] is used to prevent LT-producing enterotoxigenic *E. coli* (ETEC) diarrhea in travellers.



Infection with ETEC is the leading cause of diarrhea during travel in many developing countries, resulting in an estimated one third of cases of diarrhea in travellers to Latin America and the Caribbean, Africa and South Asia.



Populations with increased susceptibility and higher risk for more severe disease caused by ETEC include:

- People with achlorhydria (absence of gastric acid secretion)
- People with previous gastrectomy (removal of part or all of the stomach)
- People with a history of repeated severe diarrhea
- Young children aged >2 years
- People with immunodeficiency disorders

Natural immunity decreases over time. Even if you were born in a high-risk area, you should protect yourself.

ETEC diarrhea can be prevented

What can you do?

1. Get vaccinated with **DUKORAL®**
2. Watch what you eat, drink and do

Don't drink:



Tap water



Drinks with ice cubes

- Including alcoholic beverages served or shaken with ice cubes



Drinks with crushed ice

Don't eat:



Fruits that can't be peeled



Uncooked vegetables, including salad



Undercooked or raw meat or fish



Unpasteurized or unrefrigerated dairy products



Food sold by street vendors

Diarrhea caused by ETEC can easily spread.

- Wash your hands frequently
- Brush your teeth with treated or bottled water
- Avoid swimming in polluted or contaminated water
- Wipe cans and bottles before opening them



Help prevent ETEC diarrhea with
DUKORAL[®]
2 weeks before you leave for 3 months protection

Taking DUKORAL[®] before your trip

To help prevent diarrhea caused by
LT-producing enterotoxigenic *E. coli*

Store in refrigerator – DO NOT FREEZE

☐ IF YOU ARE TAKING DUKORAL[®] FOR THE FIRST TIME:



FIRST DOSE

at least 2 weeks
before your departure

Date: _____



SECOND DOSE

at least 1 week after the first dose
and at least 1 week before your
departure

Date: _____



TRIP

Protection starts 1 week
after the 2nd dose and lasts for
about 3 months

☐ IF YOU HAVE TAKEN DUKORAL[®] BEFORE AND ARE TRAVELLING AGAIN:

- If it has been less than 5 years, a single dose taken 1 week before your trip renews your protection for about 3 months
- If it has been more than 5 years since the last dose, 2 doses are needed again to renew protection

Storage and stability



STORE IN THE REFRIGERATOR at 2° to 8°C
(35° to 46°F)

DO NOT FREEZE

Do not use after expiry date.

DIN 02247208

What you need:



Spoon




Glass of water
(5 oz/150 mL)



- White sachet of powder
- Glass vial of vaccine

How to prepare DUKORAL[®]

Do not eat, drink any other liquids (including water) or take any other medication 1 hour before and 1 hour after taking DUKORAL[®]



Open the white sachet of powder and pour into 5 oz. (150 mL) of cool water. Stir gently with spoon to dissolve. Do not use any other liquid. For kids 2–6 years of age, pour away half of the powder/water mixture before Step 2.



Shake the glass vial of vaccine to mix it well.



Open the glass vial from Step 2 and pour the vaccine into the glass containing the powder/water mixture from Step 1. Stir gently with spoon to mix all ingredients.



Drink the entire contents of the glass immediately. Do not eat or drink anything, including water, for 1 hour after taking DUKORAL[®].



For a video demonstration and additional information, please visit DukoralCanada.com



Who should take DUKORAL®?

DUKORAL® is:

- An oral vaccine that is used to help prevent diarrhea caused by enterotoxigenic *E. coli* (ETEC) producing a heat-sensitive toxin (called LT) and/or cholera. The ETEC bacterium is the most common cause of diarrhea in travellers in many parts of the world.
- Used to help protect people who are travelling to an area where there is a risk of diarrhea caused by cholera and/or LT-producing ETEC. This vaccine may be given to adults and children 2 years of age and older.

Who shouldn't take DUKORAL®?

DUKORAL® is not recommended for:

- Use in pregnancy; if you are pregnant, your doctor will discuss the possible risks and benefits of having DUKORAL® during pregnancy
- Travellers experiencing acute gastrointestinal illness (e.g., diarrhea) or high temperature
- People with an allergy or hypersensitivity to any component of DUKORAL® or to formaldehyde

Talk to your doctor, nurse or pharmacist if you are not sure whether you or your child should take DUKORAL®.

Ask your healthcare professional about DUKORAL[®]



DUKORAL[®] is available at your pharmacy and travel clinic.

The cost of DUKORAL[®] may be covered by your private insurance. To find out if you are covered, contact your insurance provider and give the drug identification number (DIN) below.

DIN 02247208

The only oral vaccine for protection against enterotoxigenic *Escherichia coli* (ETEC) diarrhea, DUKORAL[®] is suitable for people as young as 2 years old. The first dose of DUKORAL[®] should be taken 2 weeks before departure, and the second should be taken 1 week before departure. DUKORAL[®] provides about 3 months of protection against enterotoxigenic *E. coli* (ETEC).

Help prevent ETEC diarrhea with
DUKORAL[®]
2 weeks before you leave for 3 months protection

Help prevent ETEC diarrhea with
DUKORAL[®]
2 weeks before you leave for 3 months protection

**WHATEVER YOU'RE PLANNING,
YOUR VACATION IS NOT A GOOD TIME FOR
ENTEROTOXIGENIC *E. COLI*/DIARRHEA.**



**Visit dukoralcanada.com for more information,
or talk to your healthcare professional.**

DUKORAL[®] does not treat diarrhea caused by enterotoxigenic *E. coli* once it develops. DUKORAL[®] will only protect against cholera and LT-producing enterotoxigenic *E. coli* (ETEC). Not everyone who gets vaccinated will be fully protected. Therefore, precautions to avoid contaminated food or water should be taken. Allergic reactions and side effects such as abdominal pain, diarrhea, fever, nausea and vomiting may occur. Rehydration measures should be taken in case of diarrhea.



VALNEVA Canada Inc.
600-3535, Saint-Charles Blvd.
Kirkland (Montreal), Quebec
H9H 5B9

© 2020 VALNEVA Canada Inc.
All trademarks used under license.
DK0132020FEB17E

