

Influenza Vaccine (surface antigen, inactivated, prepared in cell cultures)

PATIENT INFORMATION

FLUCELVAX® QUAD IS INDICATED TO BE USED IN ADULTS AND CHILDREN 6 MONTHS AND OLDER TO PREVENT INFLUENZA.

Visit flucelvax.ca for more information!

READ THIS INFORMATION CAREFULLY BEFORE YOU ARE GIVEN FLUCELVAX® QUAD.

This leaflet is a summary and will not tell you everything about this vaccine. Talk to your healthcare professional and see if **FLUCELVAX® QUAD** is right for you.

WHAT IS FLUCELVAX® QUAD USED FOR?

FLUCELVAX® QUAD is used in adults and children aged 6 months and older to prevent influenza.

Influenza, often called "the flu", is caused by infection with specific influenza viruses. New types of influenza viruses can appear each year. FLUCELVAX® QUAD vaccine contains fragments of four different types of influenza virus. Each year the World Health Organization Decides which four strains of viruses are most suitable to include in the vaccine.

You cannot catch influenza from the vaccine, as the virus in the vaccine has been killed and split into small non-infectious particles.

The National Advisory Committee on Immunization (NACI) encourages annual influenza vaccination for all Canadians who are able to have the vaccine.

Vaccination against influenza is recommended every year, for anyone wanting to lower their chance of catching influenza. FLUCELVAX® QUAD has been used by many people to lower their risk of catching the flu.



HOW DOES FLUCELVAX® QUAD WORK?

FLUCELVAX® QUAD vaccine works by helping your body protect itself against infection by the types of influenza viruses that are in the vaccine. The vaccine stimulates the body to make substances called antibodies. Antibodies fight the influenza virus. If you have been vaccinated, when you come into contact with the influenza viruses in the vaccine, your body is usually able to quickly destroy the virus, which may prevent you from getting influenza.

Your body takes a few weeks after vaccination to fully develop effective protection against the influenza virus.

Protection against influenza generally requires one dose of FLUCELVAX® QUAD vaccine. Some people including children who have not received influenza vaccination before may require two doses.

As with all vaccines, 100% protection cannot be guaranteed.

FLUCELVAX[®] QUAD is not made using eggs; there are no egg proteins in the vaccine.

The syringe and vial components do not contain latex. FLUCELVAX[®] QUAD is considered safe for use in persons with latex allergies.

FLUCELVAX® QUAD pre-filled syringes contain no preservative or antibiotics. FLUCELVAX® QUAD multi-dose vial formulation contains a preservative, but does not contain any antibiotics.

DO NOT USE FLUCELVAX® QUAD IF:

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take FLUCELVAX® QUAD. Talk about any health conditions or problems you may have, including if you or your child:

- Have or have had a reaction to vaccination with any of the following:
 - severe allergic reaction
 - difficulty breathin
 - swelling of the throa
 - fainting or collaps
 - fits or convulsions
- high temperature (greater than 38.5°C)
- severe skin reaction at the injection site, including severe bruising
- Have an infection or temperature higher than 38.5°C Your doctor may decide to delay vaccination until the illness has passed. A minor illness such as a cold is not usually a reason to delay vaccination.
- Have low immunity due to treatment with certain medicines
- Have or have had Guillain-Barré Syndrome (GBS) An illness which affects the nervous system and causes paralysis.
- Have allergies to other medicines or substances

Are pregnant or breastfeeding

Your healthcare professional will be able to discuss the potential risks and benefits of having FLUCELVAX® QUAD while you are pregnant or breastfeeding.

TELL YOUR HEALTHCARE PROFESSIONAL ABOUT ALL THE MEDICINES YOU TAKE, INCLUDING ANY DRUGS, VITAMINS, MINERALS, NATURAL SUPPLEMENTS OR ALTERNATIVE MEDICINES.



• YOUR CHILD IS UNDER 6 MONTHS OF AGE. FLUCELVAX® QUAD VACCINE IS APPROVED FOR USE IN CHILDREN AGED 6 MONTHS AND OLDER.

• YOU OR YOUR CHILD HAVE OR HAVE HAD AN ALLERGY TO FLUCELVAX® QUAD OR ANY INGREDIENT LISTED IN THIS LEAFLET.

HOW FLUCELVAX® QUAD IS GIVEN:

 $\mathsf{FLUCELVAX}^{\otimes}$ QUAD is given as an injection into a muscle, usually in the upper arm.

USUAL DOSE:

- FLUCELVAX® QUAD is given once every year as follows:
- Adults and children 9 years and over: one injection of 0.5 mL.

For children 6 months to less than 9 years old who are receiving influenza vaccine for the first time, it is recommended that a follow-up (booster) dose of FLUCELVAX® QUAD is given 4 weeks after the first dose.

If the follow-up dose is missed, talk to your healthcare professional and arrange another visit as soon as possible.

OVERDOSE:

IF YOU THINK YOU HAVE BEEN GIVEN TOO MANY DOSES OF FLUCELVAX® QUAD OR HAVE BEEN GIVEN IT BY MISTAKE, CONTACT YOUR HEALTHCARE PROFESSIONAL, HOSPITAL EMERGENCY DEPARTMENT OR REGIONAL POISON CONTROL CENTRE IMMEDIATELY, EVEN IF THERE ARE NO SYMPTOMS.

WHAT ARE POSSIBLE SIDE EFFECTS FROM USING FLUCELVAX® QUAD?

These are not all the possible side effects you may experience when taking FLUCELVAX® QUAD. If you experience any side effects not listed here, contact your healthcare professional.

The following are common or very common side effects of FLUCELVAX® QUAD. Most of these side effects are mild and do not last long. Tell your doctor if you or your child have side effects that bother you:

• Nausea, vomiting, diarrhea

• Loss of appetite

Bruising

Shivering

- Injection-site pain, reddening,
- hardening or swelling
- Headache
- Muscle or joint pain
- Tiredness



Younger children (less than 6 years of age) may experience the following common or very common side effects:

- Injection site tenderness
- Irritability
- Sleepiness
- Change of eating habits
- Fever

If you experience any of the rare **serious side effects** mentioned below, **talk to your healthcare professional**:

- Anaphylaxis: Difficulty breathing, dizziness, a weak and rapid pulse, skin rash.
- Allergic reaction: Rash, itching or hives on the skin, swelling of the face, lips, tongue, or other parts of the body.

IF YOU HAVE A TROUBLESOME SYMPTOM OR SIDE EFFECT THAT IS NOT LISTED HERE OR BECOMES BAD ENOUGH TO INTERFERE WITH YOUR DAILY ACTIVITIES, TALK TO YOUR HEALTHCARE PROFESSIONAL.



REPORTING SUSPECTED SIDE EFFECTS

Should you experience a side effect following immunization, please report it to your doctor, nurse, or pharmacist.

Should you require information related to the management of the side effect, please contact your healthcare provider. The Public Health Agency of Canada, Health Canada and Seqirus cannot provide medical advice.



If you want more information about FLUCELVAX[®] QUAD:

- Talk to your healthcare professional
- Find the full Product Monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website, the manufacturer's website (www.seqirus.ca), or by calling 1-855-358-8966.
- Visit flucelvax.ca



The Drug Identification Number (DIN) for **FLUCELVAX® QUAD** is 02494248. Your insurance company may ask for this number.

References: 1. FLUCELVAX® QUAD Product Monograph. Apr 12, 2022 2. WHO. Influenza: Recommended composition or influenza virus vaccines for use in the 2020 - 2021 Northern Hemisphere influenza season. February 28, 2020. https://www.who.int/influenza/vaccines/virus/recommendations/2020-21_north/en/ (accessed March 3, 2020).



FLUCELVAX[®] QUAD

FLUCELVAX[®] QUAD IS INDICATED TO BE USED IN ADULTS AND CHILDREN 6 MONTHS AND OLDER TO PREVENT INFLUENZA.

Visit flucelvax.ca for even more information!



Seqirus Canada Inc. Kirkland, Québec www.seqirus.ca 1.844.392.8582

Flucelvax is a registered trademark of Seqirus UK limited or its affiliates. © Seqirus Canada Inc. 2022 CAN-IV-22-0001